

WEAK IMPOSITION OF SIGNORINI BOUNDARY CONDITIONS ON THE BOUNDARY ELEMENT METHOD*

ERIK BURMAN[†], STEFAN FREI[‡], AND MATTHEW W. SCROGGS[§]

Abstract. We derive and analyze a boundary element formulation for boundary conditions involving inequalities. In particular, we focus on Signorini contact conditions. The Calderón projector is used for the system matrix, and boundary conditions are weakly imposed using a particular variational boundary operator designed using techniques from augmented Lagrangian methods. We present a complete numerical a priori error analysis and present some numerical examples to illustrate the theory.

Key words. boundary element methods, Nitsche’s method, Signorini problem, Calderón projector

AMS subject classifications. 65N38, 65R20, 74M15

DOI. 10.1137/19M1281721

1. Introduction. The application of Nitsche techniques to deal with variational inequalities has received increasing interest recently, starting from a series of works by Chouly, Hild, and Renard for elasticity problems with contact [7]. Their approach goes back to an augmented Lagrangian formulation, that was first introduced by Alart and Curnier [1].

In a previous paper [2], we have shown how Nitsche techniques can be used to impose Dirichlet, Neumann, mixed Dirichlet–Neumann, or Robin conditions weakly within boundary element methods. By using the Calderón projector, we were able to derive a unified framework that can be used for different boundary conditions.

The purpose of this article is to extend these techniques to boundary conditions involving inequalities, such as Signorini contact conditions. In particular, we consider the Laplace equation with mixed Dirichlet and Signorini boundary conditions: Find u such that

$$\begin{aligned}
 (1.1a) \quad & -\Delta u = 0 && \text{in } \Omega, \\
 (1.1b) \quad & u = g_D && \text{on } \Gamma_D, \\
 (1.1c) \quad & u \leq g_C \quad \text{and} \quad \frac{\partial u}{\partial \boldsymbol{\nu}} \leq \psi_C && \text{on } \Gamma_C, \\
 (1.1d) \quad & \left(\frac{\partial u}{\partial \boldsymbol{\nu}} - \psi_C \right) \left(u - g_C \right) = 0 && \text{on } \Gamma_C.
 \end{aligned}$$

Here $\Omega \subset \mathbb{R}^3$ denotes a polyhedral domain with outward pointing normal $\boldsymbol{\nu}$ and boundary $\Gamma := \Gamma_D \cup \Gamma_C$. We assume for simplicity that the boundary between Γ_D

*Received by the editors August 15, 2019; accepted for publication (in revised form) May 11, 2020; published electronically August 13, 2020.

<https://doi.org/10.1137/19M1281721>

Funding: The work of the first author was supported by the EPSRC grant EP/P01576X/1. The work of the second author was supported by the DFG Research Scholarship 3935/1-1.

[†]Department of Mathematics, University College London, London, WC1E 6BT, UK (e.burman@ucl.ac.uk).

[‡]Department of Mathematics and Statistics, University of Konstanz, Konstanz D-78457, Germany (stefan.frei@uni-konstanz.de).

[§]Department of Engineering, University of Cambridge, Cambridge CB2 1PZ, UK (mws48@cam.ac.uk, <http://www.msccrogs.co.uk>).

and Γ_C coincides with edges between the faces of Γ . Whenever it is ambiguous, we will write $\boldsymbol{\nu}_x$ for the outward pointing normal at the point \boldsymbol{x} . We assume that

$$g = \begin{cases} g_D & \text{in } \Gamma_D \\ g_C & \text{in } \Gamma_C \end{cases} \in L^2(\Gamma) \text{ and } \psi_C \in H^{1/2}(\Gamma_C).$$

Observe that when $\Gamma_C = \emptyset$, there exists a unique solution to (1.1) by the Lax–Milgram lemma. In the case that $\text{meas}(\Gamma_C) > 0$, the theory of Lions and Stampacchia [12] for variational inequalities yields existence and uniqueness of solutions. We assume that $u \in H^{3/2+\epsilon}(\Omega)$ for some $\epsilon > 0$.

Boundary element methods for Signorini problems were first studied by Han [11]. A variational formulation involving the Calderón projector was presented in [10]. An alternative formulation is based on Steklov–Poincaré operators [20, 22]. The numerical approaches to solve such formulations include a penalty formulation [15], operator splitting techniques [17, 23], or semi-smooth Newton methods [20, 22]. Besides the usual energy norm estimates, the latter reference includes an $L^2(\Gamma)$ -error estimate based on a duality argument. Maischak and Stephan [13] presented a posteriori error estimates and an hp -adaptive algorithm for the Signorini problem. A priori error estimates for a penalty-based hp algorithm were shown by Chernov, Maischak, and Stephan [6]. Recently, an augmented Lagrangian approach has been presented in combination with a semi-smooth Newton method [22], and variational inequalities have been successfully used for time-dependent contact problems [9].

We will consider an approach where the full Calderón projector is used and the boundary conditions are included by properly adding scaled penalty terms to the two equations. This results in formulations similar to the ones obtained for weak imposition of boundary conditions using Nitsche’s method [14]. The proposed framework is flexible and allows for the design of a range of different methods depending on the choice of weights and residuals.

An outline of this paper is as follows. In section 2, we introduce the basic boundary operators that will be needed and review some of their properties. Then, in section 3, we introduce the variational framework and review the results from [2] for the pure Dirichlet problem. In section 4, we show how the framework can be applied to Signorini boundary conditions and the mixed problem (1.1). The method is analyzed in section 5. We conclude by showing some numerical experiments in section 6.

2. Boundary operators. We define Green’s function for the Laplace operator in \mathbb{R}^3 by

$$(2.1) \quad G(\boldsymbol{x}, \boldsymbol{y}) = \frac{1}{4\pi|\boldsymbol{x} - \boldsymbol{y}|}.$$

In this paper, we focus on the problem in \mathbb{R}^3 . Similar analysis can be used for problems in \mathbb{R}^2 , in which case this definition should be replaced by $G(\boldsymbol{x}, \boldsymbol{y}) = -\log|\boldsymbol{x} - \boldsymbol{y}|/2\pi$.

In the standard fashion (see, e.g., [19, Chapter 6]), we define the single layer potential operator, $\mathcal{V} : H^{-1/2}(\Gamma) \rightarrow H^1(\Omega)$, and the double layer potential, $\mathcal{K} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$ for $v \in H^{1/2}(\Gamma)$, $\mu \in H^{-1/2}(\Gamma)$, and $\boldsymbol{x} \in \Omega \setminus \Gamma$ by

$$(2.2) \quad (\mathcal{V}\mu)(\boldsymbol{x}) := \int_{\Gamma} G(\boldsymbol{x}, \boldsymbol{y})\mu(\boldsymbol{y}) \, d\boldsymbol{y},$$

$$(2.3) \quad (\mathcal{K}v)(\boldsymbol{x}) := \int_{\Gamma} \frac{\partial G(\boldsymbol{x}, \boldsymbol{y})}{\partial \boldsymbol{\nu}_{\boldsymbol{y}}} v(\boldsymbol{y}) \, d\boldsymbol{y}.$$

We define the space $H^1(\Delta, \Omega) := \{v \in H^1(\Omega) : \Delta v \in L^2(\Omega)\}$, and the Dirichlet and Neumann traces, $\gamma_D : H^1(\Omega) \rightarrow H^{1/2}(\Gamma)$ and $\gamma_N : H^1(\Delta, \Omega) \rightarrow H^{-1/2}(\Gamma)$, by

$$(2.4) \quad \gamma_D f(\mathbf{x}) := \lim_{\Omega \ni \mathbf{y} \rightarrow \mathbf{x} \in \Gamma} f(\mathbf{y}),$$

$$(2.5) \quad \gamma_N f(\mathbf{x}) := \lim_{\Omega \ni \mathbf{y} \rightarrow \mathbf{x} \in \Gamma} \boldsymbol{\nu}_{\mathbf{x}} \cdot \nabla f(\mathbf{y}).$$

We recall that if the Dirichlet and Neumann traces of a harmonic function are known, then the potentials (2.2) and (2.3) may be used to reconstruct the function in Ω using the following relation:

$$(2.6) \quad u = -\mathcal{K}(\gamma_D u) + \mathcal{V}(\gamma_N u).$$

It is also known [19, Lemma 6.6] that for all $\mu \in H^{-1/2}(\Gamma)$, the function

$$(2.7) \quad u_\mu^\mathcal{V} := \mathcal{V}\mu$$

satisfies $-\Delta u_\mu^\mathcal{V} = 0$ and

$$(2.8) \quad \|u_\mu^\mathcal{V}\|_{H^1(\Omega)} \leq c \|\mu\|_{H^{-1/2}(\Gamma)}.$$

Similarly, in [19, Lemma 6.10] the function

$$(2.9) \quad u_v^\mathcal{K} := \mathcal{K}v$$

satisfies $-\Delta u_v^\mathcal{K} = 0$ for all $v \in H^{1/2}(\Gamma)$ and

$$(2.10) \quad \|u_v^\mathcal{K}\|_{H^1(\Omega)} \leq c \|v\|_{H^{1/2}(\Gamma)}.$$

We define $\{\gamma_D f\}_\Gamma$ and $\{\gamma_N f\}_\Gamma$ to be the averages of the interior and exterior Dirichlet and Neumann traces of f . We define the single layer, double layer, adjoint double layer, and hypersingular boundary integral operators, $\mathbf{V} : H^{-1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma)$, $\mathbf{K} : H^{1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma)$, $\mathbf{K}' : H^{-1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma)$, and $\mathbf{W} : H^{1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma)$, by

$$(2.11a) \quad (\mathbf{K}v)(\mathbf{x}) := \{\gamma_D \mathcal{K}v\}_\Gamma(\mathbf{x}), \quad (\mathbf{V}\mu)(\mathbf{x}) := \{\gamma_D \mathcal{V}\mu\}_\Gamma(\mathbf{x}),$$

$$(2.11b) \quad (\mathbf{W}v)(\mathbf{x}) := -\{\gamma_N \mathcal{K}v\}_\Gamma(\mathbf{x}), \quad (\mathbf{K}'\mu)(\mathbf{x}) := \{\gamma_N \mathcal{V}\mu\}_\Gamma(\mathbf{x}),$$

where $\mathbf{x} \in \Gamma$, $v \in H^{1/2}(\Gamma)$, and $\mu \in H^{-1/2}(\Gamma)$ [19, Chapter 6].

Next, we define the Calderón projector by

$$(2.12) \quad \mathbf{C} := \begin{pmatrix} (1 - \sigma)\text{Id} - \mathbf{K} & \mathbf{V} \\ \mathbf{W} & \sigma\text{Id} + \mathbf{K}' \end{pmatrix},$$

where σ is defined for $\mathbf{x} \in \Gamma$ by [19, equation 6.11],

$$(2.13) \quad \sigma(\mathbf{x}) = \lim_{\epsilon \rightarrow 0} \frac{1}{4\pi\epsilon^2} \int_{\mathbf{y} \in \Omega: |\mathbf{y}-\mathbf{x}|=\epsilon} d\mathbf{y}.$$

Recall that if u is a solution of (1.1), then it satisfies

$$(2.14) \quad \mathbf{C} \begin{pmatrix} \gamma_D u \\ \gamma_N u \end{pmatrix} = \begin{pmatrix} \gamma_D u \\ \gamma_N u \end{pmatrix}.$$

Taking the product of (2.14) with two test functions, and using the fact that $\sigma = \frac{1}{2}$ almost everywhere, we arrive at the following equations:

$$(2.15) \quad \langle \gamma_{\mathbb{D}} u, \mu \rangle_{\Gamma} = \langle (\tfrac{1}{2} \text{Id} - \mathbf{K}) \gamma_{\mathbb{D}} u, \mu \rangle_{\Gamma} + \langle \mathbf{V} \gamma_{\mathbb{N}} u, \mu \rangle_{\Gamma} \quad \forall \mu \in H^{-1/2}(\Gamma),$$

$$(2.16) \quad \langle \gamma_{\mathbb{N}} u, v \rangle_{\Gamma} = \langle (\tfrac{1}{2} \text{Id} + \mathbf{K}') \gamma_{\mathbb{N}} u, v \rangle_{\Gamma} + \langle \mathbf{W} \gamma_{\mathbb{D}} u, v \rangle_{\Gamma} \quad \forall v \in H^{1/2}(\Gamma).$$

For a more compact notation, we introduce $\lambda = \gamma_{\mathbb{N}} u$, $u = \gamma_{\mathbb{D}} u$, and the Calderón form

$$(2.17) \quad \mathcal{C}[(u, \lambda), (v, \mu)] := \langle (\tfrac{1}{2} \text{Id} - \mathbf{K}) u, \mu \rangle_{\Gamma} + \langle \mathbf{V} \lambda, \mu \rangle_{\Gamma} \\ + \langle (\tfrac{1}{2} \text{Id} + \mathbf{K}') \lambda, v \rangle_{\Gamma} + \langle \mathbf{W} u, v \rangle_{\Gamma}.$$

We may then rewrite (2.15) and (2.16) as

$$(2.18) \quad \mathcal{C}[(u, \lambda), (v, \mu)] = \langle u, \mu \rangle_{\Gamma} + \langle \lambda, v \rangle_{\Gamma}.$$

We will also frequently use the multitrace form, defined by

$$(2.19) \quad \mathcal{A}[(u, \lambda), (v, \mu)] := -\langle \mathbf{K} u, \mu \rangle_{\Gamma} + \langle \mathbf{V} \lambda, \mu \rangle_{\Gamma} + \langle \mathbf{K}' \lambda, v \rangle_{\Gamma} + \langle \mathbf{W} u, v \rangle_{\Gamma}.$$

Using this, we may rewrite (2.18) as

$$(2.20) \quad \mathcal{A}[(u, \lambda), (v, \mu)] = \tfrac{1}{2} \langle u, \mu \rangle_{\Gamma} + \tfrac{1}{2} \langle \lambda, v \rangle_{\Gamma}.$$

To quantify the two traces, we introduce the product space

$$\mathbb{V} := H^{1/2}(\Gamma) \times H^{-1/2}(\Gamma)$$

and the associated norm

$$\|(v, \mu)\|_{\mathbb{V}} := \|v\|_{H^{1/2}(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)}.$$

The continuity and coercivity of \mathcal{A} are immediate consequences of the properties of the operators \mathbf{V} , \mathbf{K} , \mathbf{K}' , and \mathbf{W} :

LEMMA 2.1 (continuity & coercivity). *There exists $C > 0$ such that*

$$|\mathcal{A}[(w, \eta), (v, \mu)]| \leq C \|(w, \eta)\|_{\mathbb{V}} \|(v, \mu)\|_{\mathbb{V}} \quad \forall (w, \eta), (v, \mu) \in \mathbb{V}.$$

There exists $\alpha > 0$ such that

$$\alpha \left(\|v\|_{H^{1/2}(\Gamma)}^2 + \|\mu\|_{H^{-1/2}(\Gamma)}^2 \right) \leq \mathcal{A}[(v, \mu), (v, \mu)] \quad \forall (v, \mu) \in \mathbb{V}.$$

Proof. See [2] for the proof. \square

3. Discretization and weak imposition of Dirichlet boundary conditions. In this section, we introduce the discrete spaces and review briefly how (non-homogeneous) Dirichlet boundary conditions can be imposed weakly within the variational formulations introduced above. For a detailed derivation, and for different boundary conditions, we refer to [2].

To reduce the number of constants that appear, we introduce the following notation:

- If $\exists C > 0$ such that $a \leq Cb$, then we write $a \lesssim b$.




FIG. 1. A grid (left), the barycentric refinement of the grid (center), and the dual grid (right). In a typical example, the initial grid will not be flat, and so the elements of the dual grid will not necessarily be flat.

- If $a \lesssim b$ and $b \lesssim a$, then we write $a \approx b$.

We assume that Ω is a polygonal domain with faces denoted by $\{\Gamma_i\}_{i=1}^M$. We introduce a family of conforming, shape regular triangulations of Γ , $\{\mathcal{T}_h\}_{h>0}$, indexed by the largest element diameter of the mesh, h . We let $T_1, \dots, T_m \in \mathcal{T}_h$ be the triangles of a triangulation.

We consider the following finite element spaces:

$$\begin{aligned} P_h^k(\Gamma) &:= \{v_h \in C^0(\Gamma) : v_h|_{T_i} \in \mathbb{P}_k(T_i) \text{ for every } T_i \in \mathcal{T}_h\}, \\ DP_h^l(\Gamma) &:= \{v_h \in L^2(\Gamma) : v_h|_{T_i} \in \mathbb{P}_l(T_i) \text{ for every } T_i \in \mathcal{T}_h\}, \\ \widetilde{DP}_h^l(\Gamma) &:= \{v_h \in DP_h^l(\Gamma) : v_h|_{\Gamma_i} \in C^0(\Gamma_i) \text{ for } i = 1, \dots, M\}, \end{aligned}$$

where $\mathbb{P}_k(T_i)$ denotes the space of polynomials of order less than or equal to k on the triangle T_i .

In addition, we consider the space $\text{DUAL}_h^0(\Gamma)$ of piecewise constant functions on the barycentric dual grid, as shown in Figure 1. On nonsmooth domains, these spaces have lower order approximation properties than the standard space $DP_h^0(\Gamma)$, as given in the following lemma.

LEMMA 3.1. *Let $\mu \in H^s(\Gamma)$. If Γ consists of a finite number of smooth faces meeting at edges, then*

$$\inf_{\eta_h \in \text{DUAL}_h^0(\Gamma)} \|\mu - \eta_h\|_{H^{-1/2}(\Gamma)} \lesssim h^{\xi+1/2} \|\mu\|_{H^\xi(\Gamma)},$$

where $\xi = \min(\frac{1}{2}, s)$. If Γ is smooth, then the same result holds with $\xi = \min(1, s)$.

Proof. See [16, Appendix 2] for the proof. \square

We observe that $P_h^k(\Gamma) \subset H^{1/2}(\Gamma)$, $DP_h^l(\Gamma) \subset L^2(\Gamma)$, $\widetilde{DP}_h^l(\Gamma) \subset L^2(\Gamma)$, and $\text{DUAL}_h^0(\Gamma) \subset L^2(\Gamma)$. We define the discrete product space

$$\mathbb{V}_h := P_h^k(\Gamma) \times \Lambda_h^l,$$

where Λ_h^l can be any of the spaces $DP_h^l(\Gamma)$, $\widetilde{DP}_h^l(\Gamma)$, or $\text{DUAL}_h^0(\Gamma)$.

3.1. Dirichlet boundary conditions. Let us, for the moment, assume that $\Gamma \equiv \Gamma_D$. Then, the basic idea is to add the following suitably weighted boundary residual to the weak formulation:

$$(3.1) \quad R_{\Gamma_D}(u_h, \lambda_h) := \beta_D^{1/2} (g_D - u_h).$$

This is defined such that $R_{\Gamma_D}(u_h, \lambda_h) = 0$ is equivalent to the boundary condition (1.1b). We obtain an expression of the form

$$(3.2) \quad \mathcal{C}[(u_h, \lambda_h), (v_h, \mu_h)] = \langle u_h, \mu_h \rangle_{\Gamma} + \langle \lambda_h, v_h \rangle_{\Gamma} + \langle R_{\Gamma_D}(u_h, \lambda_h), \beta_1 v_h + \beta_2 \mu_h \rangle_{\Gamma},$$

or equivalently,

$$(3.3) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] = \frac{1}{2} \langle u_h, \mu_h \rangle_{\Gamma} + \frac{1}{2} \langle \lambda_h, v_h \rangle_{\Gamma} + \langle R_{\Gamma_D}(u_h, \lambda_h), \beta_1 v_h + \beta_2 \mu_h \rangle_{\Gamma},$$

where β_1 and β_2 are problem dependent scaling operators that can be chosen as a function of the physical parameters in order to obtain robustness of the method.

For the Dirichlet problem, we choose $\beta_1 = \beta_D^{1/2}$, $\beta_2 = \beta_D^{-1/2}$, where different choices for β_D in the range $0 \leq \beta_D \lesssim h^{-1}$ are possible. Inserting this into (3.3), we obtain the formulation

$$(3.4) \quad \mathcal{A}[(u, \lambda), (v_h, \mu_h)] - \frac{1}{2} \langle \lambda_h, v_h \rangle_{\Gamma_D} + \frac{1}{2} \langle u_h, \mu_h \rangle_{\Gamma_D} + \langle \beta_D u_h, v_h \rangle_{\Gamma_D} \\ = \langle g_D, \beta_D v_h + \mu_h \rangle_{\Gamma_D}.$$

By formally identifying λ_h with $\partial_\nu u_h$ and μ_h with $\partial_\nu v_h$, we obtain the classical (non-symmetric) Nitsche's method (up to the multiplicative factor $\frac{1}{2}$).

For a more compact notation, we introduce the boundary operator associated with the nonhomogeneous Dirichlet condition

$$(3.5) \quad \mathcal{B}_D[(u_h, \lambda_h), (v_h, \mu_h)] := -\frac{1}{2} \langle \lambda_h, v_h \rangle_{\Gamma_D} + \frac{1}{2} \langle u_h, \mu_h \rangle_{\Gamma_D} + \langle \beta_D u_h, v_h \rangle_{\Gamma_D},$$

the operator corresponding to the left-hand side

$$(3.6) \quad \mathcal{A}_D[(u_h, \lambda_h), (v_h, \mu_h)] := \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_D[(u_h, \lambda_h), (v_h, \mu_h)]$$

and the operator associated with the right-hand side

$$(3.7) \quad \mathcal{L}_D(v_h, \mu_h) := \langle g_D, \beta_D v_h + \mu_h \rangle_{\Gamma_D}.$$

Using these and (3.4), we arrive at the following boundary element formulation: Find $(u_h, \lambda_h) \in \mathbb{V}_h$ such that

$$(3.8) \quad \mathcal{A}_D[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}_D(v_h, \mu_h) \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

We introduce the following \mathcal{B}_D -norm:

$$\|(v, \mu)\|_{\mathcal{B}_D} := \|(v, \mu)\|_{\mathbb{V}} + \beta_D^{1/2} \|v\|_{\Gamma_D},$$

and summarize the properties of the bilinear form \mathcal{A}_D in the following lemma.

LEMMA 3.2 (properties of the bilinear form). *Let \mathbb{W} be a product Hilbert space for the primal and flux variables, such that $\mathbb{V} \subset \mathbb{W}$. The bilinear form has the following properties:*

PROPERTY 1 (coercivity). *If $\beta_D = 0$ or if there exists $\beta_{\min} > 0$ (independent of h) such that $\beta_D > \beta_{\min}$, then there exists $\alpha > 0$ such that $\forall (v, \mu) \in \mathbb{W}$,*

$$\alpha \|(v, \mu)\|_{\mathcal{B}_D} \leq \mathcal{A}_D[(v, \mu), (v, \mu)].$$

PROPERTY 2 (continuity). *There exists $M > 0$ such that $\forall (w, \eta), (v, \mu) \in \mathbb{W}$,*

$$|\mathcal{A}_D[(v, \mu), (w, \eta)]| \leq M \|(v, \mu)\|_{\mathcal{B}_D} \|(w, \eta)\|_{\mathcal{B}_D}.$$

Proof. See [2, section 4.1] for the proof. □

4. Weak imposition of Signorini boundary conditions. Recently, Chouly, Hild, and Renard [7, 8] showed how contact problems can be treated in the context of Nitsche's method. We will show here how we may use arguments similar to theirs in the present framework to integrate unilateral contact seamlessly. The result is a nonlinear system to which one may apply Newton's method or a fixed-point iteration in a straightforward manner. We prove existence and uniqueness of solutions to the nonlinear system and optimal order error estimates.

For the derivation of the formulation on the contact boundary we will first omit the Dirichlet part, letting $\Gamma = \Gamma_C$. To impose the contact conditions, we recall the following relations, introduced by Alart and Curnier [1], with $[x]_{\pm} := \pm \max(0, \pm x)$:

$$(4.1) \quad (u - g_C) = [(u - g_C) - \tau^{-1}(\lambda - \psi_C)]_- \quad \text{on } \Gamma_C,$$

$$(4.2) \quad (\lambda - \psi_C) = -[\tau(u - g_C) - (\lambda - \psi_C)]_+ \quad \text{on } \Gamma_C$$

for all $\tau > 0$. It is straightforward [7] to show that each of these two conditions is equivalent to the contact boundary conditions (1.1c) and (1.1d).

To simplify the notation, we introduce the operators

$$P^\tau(u_h, \lambda_h) := \tau(u_h - g_C) - (\lambda_h - \psi_C) \quad \text{and} \quad P_0^\tau(u_h, \lambda_h) := \tau u_h - \lambda_h.$$

Using (4.1), we arrive at the following boundary term for the contact conditions:

$$(4.3) \quad R_{\Gamma_C}^1(u_h, \lambda_h) = (g_C - u_h) + \tau^{-1} [P^\tau(u_h, \lambda_h)]_-.$$

Alternatively, by using (4.2), we arrive at the following boundary term:

$$(4.4) \quad R_{\Gamma_C}^2(u_h, \lambda_h) = \tau^{-1} ((\psi_C - \lambda_h) - [P^\tau(u_h, \lambda_h)]_+).$$

By using the fact that $x = [x]_+ + [x]_-$, it can be shown that (4.3) and (4.4) are equal.

Substituting (4.3) into (3.3) and using the weights $\beta_1 = \tau$ and $\beta_2 = 1$, we obtain

$$(4.5) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \frac{1}{2} \langle \mu_h, u_h \rangle_{\Gamma_C} + \langle \tau u_h - \frac{1}{2} \lambda_h, v_h \rangle_{\Gamma_C} \\ - \langle [P^\tau(u_h, \lambda_h)]_-, v_h + \tau^{-1} \mu_h \rangle_{\Gamma_C} = \langle g_C, \tau v_h + \mu_h \rangle_{\Gamma_C}.$$

Using (4.4), we have

$$(4.6) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \frac{1}{2} \langle \lambda_h, v_h \rangle_{\Gamma_C} + \langle \tau^{-1} \lambda_h - \frac{1}{2} u_h, \mu_h \rangle_{\Gamma_C} \\ + \langle [P^\tau(u_h, \lambda_h)]_+, v_h + \tau^{-1} \mu_h \rangle_{\Gamma_C} = \langle \psi_C, v_h + \tau^{-1} \mu_h \rangle_{\Gamma_C}.$$

We see that (4.6) is similar to the nonsymmetric version of the method proposed in [8] and (4.5) is similar to the nonsymmetric Nitsche formulation for contact discussed in [5]. As pointed out in the latter reference, the two formulations are equivalent, with the same solutions. In what follows, we focus exclusively on the variant (4.6).

Defining

$$(4.7) \quad \mathcal{B}_C[(u_h, \lambda_h), (v_h, \mu_h)] := \frac{1}{2} \langle \lambda_h, v_h \rangle_{\Gamma_C} + \langle \tau^{-1} \lambda_h - \frac{1}{2} u_h, \mu_h \rangle_{\Gamma_C} \\ + \langle [P^\tau(u_h, \lambda_h)]_+, v_h + \tau^{-1} \mu_h \rangle_{\Gamma_C},$$

$$(4.8) \quad \mathcal{L}_C(v_h, \mu_h) := \langle \psi_C, v_h + \tau^{-1} \mu_h \rangle_{\Gamma_C},$$

$$(4.9) \quad \mathcal{A}_C[(u_h, \lambda_h), (v_h, \mu_h)] := \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_C[(u_h, \lambda_h), (v_h, \mu_h)],$$

we arrive at the boundary element method formulation: Find $(u_h, \lambda_h) \in \mathbb{V}_h$ such that

$$(4.10) \quad \mathcal{A}_C[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}_C(v_h, \mu_h) \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

4.1. Mixed Dirichlet and contact boundary conditions. Combining the formulations for the Dirichlet and contact conditions, we arrive at the following boundary element method for the problem (1.1): Find $(u_h, \lambda_h) \in \mathbb{V}_h$ such that

$$(4.11) \quad \mathcal{A}_D[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_C[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}_D(v_h, \mu_h) + \mathcal{L}_C(v_h, \mu_h) \\ \forall (v_h, \mu_h) \in \mathbb{V}_h,$$

where \mathcal{A}_D , \mathcal{L}_D , \mathcal{B}_C , and \mathcal{L}_C are defined in (3.6), (3.7), (4.7), and (4.8). For discretization, we use the assumptions and spaces introduced in section 3. Note that the formulation (4.11) is consistent, i.e., the continuous solution (u, λ) to (1.1) fulfills (4.11) for all $(v_h, \mu_h) \in \mathbb{V}_h$.

5. Analysis. In this section, we prove the existence of unique solutions to the nonlinear system of equations (4.11) as well as optimal error estimates.

We assume that the solution (u, λ) of (1.1) lies in $\mathbb{W} := H^{1+\epsilon}(\Gamma) \times H^\epsilon(\tilde{\Gamma})$ for some $\epsilon \in (0, 1/2]$, where $\tilde{\Gamma} = \cup_{i=1}^M \Gamma_i \setminus \partial\Gamma_i$ is the set of boundary points that lie in the interior of the faces Γ_i . As the normal vectors ν_x are discontinuous between faces, we cannot expect a higher global regularity for λ .

We define the distance function d_C and norm $\|\cdot\|_*$ for $(v, \mu), (w, \eta) \in \mathbb{W}$, by

$$(5.1) \quad d_C((v, \mu), (w, \eta)) := \|(v - w, \mu - \eta)\|_{\mathcal{B}_D} \\ + \|\tau^{-\frac{1}{2}}(\mu - \eta + [P^\tau(v, \mu)]_+ - [P^\tau(w, \eta)]_+)\|_{\Gamma_C},$$

$$(5.2) \quad \|(v, \mu)\|_* := \|(v, \mu)\|_{\mathcal{B}_D} + \|\tau^{\frac{1}{2}}v\|_{\Gamma_C} + \|\tau^{-\frac{1}{2}}\mu\|_{\Gamma_C}.$$

We note that due to the appearance of $[\cdot]_+$ in its second term, d_C is not a norm. d_C does provide a bound on the error; however, as for all $(v, \mu) \in \mathbb{W}$, $d_C((v, \mu), (0, 0)) \geq \|(v, \mu)\|_{\mathcal{B}_D} \geq \|(v, \mu)\|_*$.

When proving this section's results, we will use properties of the $[\cdot]_+$ function that are given in the following lemma.

LEMMA 5.1. For all $a, b \in \mathbb{R}$,

$$(5.3) \quad ([a]_+ - [b]_+)^2 \leq ([a]_+ - [b]_+)(a - b),$$

$$(5.4) \quad |[a]_+ - [b]_+| \leq |a - b|.$$

Proof. For a proof of these well-known properties, see, e.g., [7]. \square

We now prove a result analogous to the coercivity assumption in [2].

LEMMA 5.2. If there is $\beta_{\min} > 0$, independent of h , such that $\beta_D > \beta_{\min}$, then there is $\alpha > 0$ such that for all $(v, \mu), (w, \eta) \in \mathbb{W}$,

$$\alpha (d_C((v, \mu), (w, \eta)))^2 \leq (\mathcal{A} + \mathcal{B}_D)[(v - w, \mu - \eta), (v - w, \mu - \eta)] \\ + \mathcal{B}_C[(v, \mu), (v - w, \mu - \eta)] - \mathcal{B}_C[(w, \eta), (v - w, \mu - \eta)].$$

Proof. From the analysis of the Dirichlet problem (Lemma 3.2) we know that when $\beta_D > \beta_{\min} > 0$,

$$(5.5) \quad \alpha \|(v - w, \mu - \eta)\|_{\mathcal{B}_D}^2 \leq (\mathcal{A} + \mathcal{B}_D)[(v - w, \mu - \eta), (v - w, \mu - \eta)].$$

Introducing the notation $\delta P := [P^\tau(v, \mu)]_+ - [P^\tau(w, \eta)]_+$, we have

$$(5.6) \quad \mathcal{B}_C[(v, \mu), (v - w, \mu - \eta)] - \mathcal{B}_C[(w, \eta), (v - w, \mu - \eta)] \\ = \tau^{-1} \|\mu - \eta\|_{\Gamma_C}^2 + \langle \delta P, v - w + \tau^{-1}(\mu - \eta) \rangle_{\Gamma_C}.$$

To estimate the expression on the right-hand side, we use

$$\tau^{-1} \|\mu - \eta + \delta P\|_{\Gamma_C}^2 = \tau^{-1} (\|\mu - \eta\|_{\Gamma_C}^2 + \|\delta P\|_{\Gamma_C}^2 + 2 \langle \mu - \eta, \delta P \rangle_{\Gamma_C}).$$

Using (5.3), this implies the bound

$$\begin{aligned} \tau^{-1} \|\mu - \eta + \delta P\|_{\Gamma_C}^2 &\leq \tau^{-1} (\|\mu - \eta\|_{\Gamma_C}^2 + \langle \delta P, P_0^\tau(v - w, \mu - \eta) \rangle_{\Gamma_C} + 2 \langle \mu - \eta, \delta P \rangle_{\Gamma_C}). \end{aligned}$$

Observing that $P_0^\tau(v - w, \mu - \eta) + 2(\mu - \eta) = \tau(v - w) + \mu - \eta$, we infer that

$$(5.7) \quad \tau^{-1} \|\mu - \eta + \delta P\|_{\Gamma_C}^2 \leq \mathcal{B}_C[(v, \mu), (v - w, \mu - \eta)] - \mathcal{B}_C[(w, \eta), (v - w, \mu - \eta)].$$

We conclude the proof by noting that

$$\begin{aligned} (d_C((v, \mu), (w, \eta)))^2 &\lesssim \|(v - w, \mu - \eta)\|_{\mathcal{B}_D}^2 \\ &\quad + \tau^{-1} \|\mu - \eta + [P^\tau(v, \mu)]_+ - [P^\tau(w, \eta)]_+\|_{\Gamma_C}^2, \end{aligned}$$

and applying (5.5) and (5.7). \square

Next, we prove a result analogous to the discrete coercivity assumption in [2].

LEMMA 5.3. *If there is $\beta_{\min} > 0$, independent of h , such that $\beta_D > \beta_{\min}$, then there is $\alpha > 0$ such that for all $(v_h, \mu_h) \in \mathbb{V}_h$,*

$$\begin{aligned} \alpha \left(\|(v_h, \mu_h)\|_{\mathcal{B}_D} + \|\tau^{-\frac{1}{2}} (\mu_h + [P^\tau(v_h, \mu_h)]_+)\|_{\Gamma_C} \right)^2 \\ \leq (\mathcal{A} + \mathcal{B}_D + \mathcal{B}_C)[(v_h, \mu_h), (v_h, \mu_h)] - \langle [P^\tau(v_h, \mu_h)]_+, g_C - \tau^{-1} \psi_C \rangle_{\Gamma_C}. \end{aligned}$$

Proof. The proof is similar to that of Lemma 5.2, but with μ_h and v_h instead of $\mu - \eta$ and $v - w$. The appearance of the data term in the right-hand side is due to the relation

$$\begin{aligned} \tau^{-1} \|[P^\tau(v_h, \mu_h)]_+\|_{\Gamma_C}^2 + 2\tau^{-1} \langle \mu_h, [P^\tau(v_h, \mu_h)]_+ \rangle_{\Gamma_C} + \tau^{-1} \|\mu_h\|_{\Gamma_C}^2 \\ = \tau^{-1} \langle [P^\tau(v_h, \mu_h)]_+, P^\tau(v_h, \mu_h) \rangle_{\Gamma_C} + \tau^{-1} \|\mu_h\|_{\Gamma_C}^2 \\ = \langle [P^\tau(v_h, \mu_h)]_+, u_h + \tau^{-1} \mu_h \rangle_{\Gamma_C} \\ \quad - \langle [P^\tau(v_h, \mu_h)]_+, g_C - \tau^{-1} \psi_C \rangle_{\Gamma_C} + \tau^{-1} \|\mu_h\|_{\Gamma_C}^2 \\ = \mathcal{B}_C[(v_h, \mu_h), (v_h, \mu_h)] - \langle [P^\tau(v_h, \mu_h)]_+, g_C - \tau^{-1} \psi_C \rangle_{\Gamma_C}. \quad \square \end{aligned}$$

Using Lemmas 5.2 and 5.3, we may now prove that (4.11) is well-posed.

THEOREM 5.4. *The finite dimensional nonlinear system (4.11) admits a unique solution.*

Proof. To prove the existence of a solution, we show the continuity and the positivity of the nonlinear operator $\mathcal{A} + \mathcal{B}_D + \mathcal{B}_C$. This allows us to apply Brouwer's fixed point theorem; see, e.g., [21, Chapter 2, Lemma 1.4].

We define $F : \mathbb{V}_h \rightarrow \mathbb{V}_h$ for $(v_h, \mu_h) \in \mathbb{V}_h$, by

$$\begin{aligned} (F(v_h, \mu_h), (w_h, \eta_h))_\Gamma &= (\mathcal{A} + \mathcal{B}_D + \mathcal{B}_C)[(v_h, \mu_h), (w_h, \eta_h)] \\ &\quad - \mathcal{L}_D(w_h, \eta_h) - \mathcal{L}_C(w_h, \eta_h) \end{aligned}$$

for all $(w_h, \eta_h) \in \mathbb{V}_h$. We may write the nonlinear system (4.11) as

$$(5.8) \quad \langle \mathbf{F}(v_h, \mu_h), (w_h, \eta_h) \rangle_\Gamma = 0 \quad \forall (w_h, \eta_h) \in \mathbb{V}_h.$$

For fixed h , by the equivalence of norms on discrete spaces, there exist $c_1, c_2 > 0$ such that for all $(v_h, \mu_h) \in \mathbb{V}_h$,

$$c_1 \|(v_h, \mu_h)\|_\Gamma \leq \|(v_h, \mu_h)\|_{\mathcal{B}_D} \leq c_2 \|(v_h, \mu_h)\|_\Gamma.$$

To show positivity, we let $(v_h, \mu_h) \in \mathbb{V}_h$. Using Lemma 5.3, we see that

$$\begin{aligned} \langle \mathbf{F}(v_h, \mu_h), (v_h, \mu_h) \rangle_\Gamma &\geq \alpha \|(v_h, \mu_h)\|_{\mathcal{B}_D}^2 + \alpha \tau^{-1} \|\mu_h + [P^\tau(v_h, \mu_h)]_+\|_{\Gamma_C}^2 \\ &\quad + \langle [P^\tau(v_h, \mu_h)]_+, g_C - \tau^{-1} \psi_C \rangle_{\Gamma_C} - \mathcal{L}_D(v_h, \mu_h) - \mathcal{L}_C(v_h, \mu_h). \end{aligned}$$

Using the Cauchy–Schwarz inequality and an arithmetic-geometric inequality, we see that there exists $C_{g_C \psi_C} > 0$ such that

$$\begin{aligned} &\langle [P^\tau(v_h, \mu_h)]_+, g_C - \tau^{-1} \psi_C \rangle_{\Gamma_C} - \mathcal{L}_D(v_h, \mu_h) - \mathcal{L}_C(v_h, \mu_h) \\ &= \langle [P^\tau(v_h, \mu_h)]_+ + \mu_h, g_C - \tau^{-1} \psi_C \rangle_{\Gamma_C} - \langle \mu_h, g_C - \tau^{-1} \psi_C \rangle_{\Gamma_C} \\ &\quad - \langle g_D, \beta_D v_h + \mu_h \rangle_{\Gamma_D} - \langle \psi_C, v_h + \tau^{-1} \mu_h \rangle_{\Gamma_C} \\ &\geq -C_{g_C \psi_C}^2 - \frac{\alpha}{2} (\|(v_h, \mu_h)\|_{\mathcal{B}_D}^2 + \tau^{-1} \|\mu_h + [P^\tau(v_h, \mu_h)]_+\|_{\Gamma_C}^2). \end{aligned}$$

Using norm equivalence, we obtain

$$\begin{aligned} &\langle \mathbf{F}(v_h, \mu_h), (v_h, \mu_h) \rangle_\Gamma \\ &\geq \frac{\alpha}{2} (\|(v_h, \mu_h)\|_{\mathcal{B}_D}^2 + \tau^{-1} \|\mu_h + [P^\tau(v_h, \mu_h)]_+\|_{\Gamma_C}^2) - C_{g_C \psi_C}^2 \\ &\geq C' \|(v_h, \mu_h)\|_\Gamma^2 - C_{g_C \psi_C}^2 \end{aligned}$$

for some $C' > 0$. We conclude that for all $(v_h, \mu_h) \in \mathbb{V}_h$ with

$$\|(v_h, \mu_h)\|_\Gamma^2 > \frac{C_{g_C \psi_C}^2}{C'} + 1,$$

there holds $\langle \mathbf{F}(v_h, \mu_h), (v_h, \mu_h) \rangle_\Gamma > 0$.

To show continuity, let $(v_h^1, \mu_h^1), (v_h^2, \mu_h^2) \in \mathbb{V}_h$. We have for all $(w_h, \eta_h) \in \mathbb{V}_h$,

$$\begin{aligned} &\langle \mathbf{F}(v_h^1, \mu_h^1) - \mathbf{F}(v_h^2, \mu_h^2), (w_h, \eta_h) \rangle_\Gamma \\ &= \left\langle [P^\tau(v_h^1, \mu_h^1)]_+ - [P^\tau(v_h^2, \mu_h^2)]_+, w_h + \tau^{-1} \eta_h \right\rangle_{\Gamma_C} \\ &\quad + \frac{1}{2} \langle \mu_h^1 - \mu_h^2, w_h + \tau^{-1} \eta_h \rangle_\Gamma - \frac{1}{2} \langle v_h^1 - v_h^2, \mu_h^1 - \mu_h^2 \rangle_{\Gamma_C} \\ &\quad + (\mathcal{A} + \mathcal{B}_D)[(v_h^1 - v_h^2, \mu_h^1 - \mu_h^2), (w_h, \eta_h)] \\ &\leq (\tau \|v_h^1 - v_h^2\|_{\Gamma_C} + \|\mu_h^1 - \mu_h^2\|_{\Gamma_C}) (\|w_h\|_{\Gamma_C} + \tau^{-1} \|\eta_h\|_{\Gamma_C}), \end{aligned}$$

where we have used (5.4). By norm equivalence, this means that

$$\frac{\langle \mathbf{F}(v_h^1, \mu_h^1) - \mathbf{F}(v_h^2, \mu_h^2), (w_h, \eta_h) \rangle_\Gamma}{\|(w_h, \eta_h)\|_\Gamma} \leq C \|(v_h^1 - v_h^2, \mu_h^1 - \mu_h^2)\|_\Gamma$$

showing that \mathbf{F} is continuous.

It then follows by Brouwer’s fixed point theorem [21, Chapter 2, Lemma 1.4] that there exists a solution to (5.8) and hence also to (4.11).

Uniqueness is an immediate consequence of Lemma 5.2. Assume that (u_h^1, λ_h^1) and (u_h^2, λ_h^2) are solutions to (4.11). We immediately see that

$$\alpha (d_C ((u_h^1, \lambda_h^1), (u_h^2, \lambda_h^2)))^2 = 0,$$

and we conclude that the solution is unique. □

We now proceed to prove the following best approximation result.

LEMMA 5.5. *Let $(u, \lambda) \in \mathbb{W}$ be the solution of (1.1) and $(u_h, \lambda_h) \in \mathbb{V}_h$ the solution of (4.11). Then there holds*

$$d_C ((u, \lambda), (u_h, \lambda_h)) \leq C \inf_{(v_h, \mu_h) \in \mathbb{V}_h} \|(u - v_h, \lambda - \mu_h)\|_*.$$

Proof. Using Lemma 5.2 and Galerkin orthogonality, we see that for arbitrary $(v_h, \mu_h) \in \mathbb{V}_h$,

$$\begin{aligned} & \alpha (d_C ((u, \lambda), (u_h, \lambda_h)))^2 \\ & \leq (\mathcal{A} + \mathcal{B}_D)[(u - u_h, \lambda - \lambda_h), (u - u_h, \lambda - \lambda_h)] \\ & \quad + \mathcal{B}_C[(u, \lambda), (u - u_h, \lambda - \lambda_h)] - \mathcal{B}_C[(u_h, \lambda_h), (u - u_h, \lambda - \lambda_h)] \\ & = (\mathcal{A} + \mathcal{B}_D)[(u - u_h, \lambda - \lambda_h), (u - v_h, \lambda - \mu_h)] \\ & \quad + \mathcal{B}_C[(u, \lambda), (u - v_h, \lambda - \mu_h)] - \mathcal{B}_C[(u_h, \lambda_h), (u - v_h, \lambda - \mu_h)]. \end{aligned}$$

Next, we use

$$\begin{aligned} & \mathcal{B}_C[(u, \lambda), (u - v_h, \lambda - \mu_h)] - \mathcal{B}_C[(u_h, \lambda_h), (u - v_h, \lambda - \mu_h)] \\ & = \langle \lambda - \lambda_h + [P^\tau(u, \lambda)]_+ - [P^\tau(u_h, \lambda_h)]_+, (u - v_h) + \tau^{-1}(\lambda - \mu_h) \rangle_{\Gamma_C} \\ & \quad - \frac{1}{2} \langle u - u_h, \lambda - \mu_h \rangle_{\Gamma_C} - \frac{1}{2} \langle \lambda - \lambda_h, u - v_h \rangle_{\Gamma_C} \end{aligned}$$

to show that

$$\begin{aligned} & (\mathcal{A} + \mathcal{B}_D)[(u - u_h, \lambda - \lambda_h), (u - u_h, \lambda - \lambda_h)] \\ & \quad + \mathcal{B}_C[(u, \lambda), (u - u_h, \lambda - \lambda_h)] - \mathcal{B}_C[(u_h, \lambda_h), (u - u_h, \lambda - \lambda_h)] \\ & = \underbrace{(\mathcal{A} + \mathcal{B}_D)[(u - u_h, \lambda - \lambda_h), (u - v_h, \lambda - \mu_h)]}_{(I)} \\ & \quad - \underbrace{\frac{1}{2} \langle u - u_h, \lambda - \mu_h \rangle_{\Gamma_C} - \frac{1}{2} \langle \lambda - \lambda_h, u - v_h \rangle_{\Gamma_C}}_{(II)} \\ & \quad + \underbrace{\langle \lambda - \lambda_h + [P^\tau(u, \lambda)]_+ - [P^\tau(u_h, \lambda_h)]_+, (u - v_h) + \tau^{-1}(\lambda - \mu_h) \rangle_{\Gamma_C}}_{(III)}. \end{aligned}$$

We estimate the three parts of the right-hand side separately. For the first term, we use the continuity of $\mathcal{A} + \mathcal{B}_D$ (Lemma 3.2) to obtain

$$(I) \leq M \|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_D} \|(u - v_h, \lambda - \mu_h)\|_{\mathcal{B}_D}.$$

For the second line, we use $H^{1/2}(\Gamma)$ – $H^{-1/2}(\Gamma)$ duality and the Cauchy–Schwarz inequality to obtain

$$(II) \leq \|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_D} \|(u - v_h, \lambda - \mu_h)\|_{\mathcal{B}_D}.$$

For the last term, we use the Cauchy–Schwarz inequality to get

$$\begin{aligned} \text{(III)} \leq & \|\tau^{-1/2}(\lambda - \lambda_h + [P^\tau(u, \lambda)]_+ - [P^\tau(u_h, \lambda_h)]_+)\|_{\Gamma_C} \\ & \cdot \left(\|\tau^{1/2}(u - v_h)\|_{\Gamma_C} + \|\tau^{-1/2}(\lambda - \mu_h)\|_{\Gamma_C} \right). \end{aligned}$$

Collecting these bounds, we see that

$$d_C((u, \lambda), (u_h, \lambda_h))^2 \lesssim d_C((u, \lambda), (u_h, \lambda_h)) \|(u - v_h, \lambda - \mu_h)\|_*.$$

Dividing through by $d_C((u, \lambda), (u_h, \lambda_h))$ and taking the infimum yields the desired result. \square

We now prove the main result of this section, an a priori bound on the error of the solution of (4.11).

THEOREM 5.6. *Let $(u, \lambda) \in H^s(\Gamma) \times H^r(\bar{\Gamma})$ for some $s \geq 1, r \geq 0$, and let $(u_h, \lambda_h) \in P_h^k(\Gamma) \times \Lambda_h^l$ be the solutions of (1.1) and the discrete problem (4.11), respectively. If there is $\beta_{\min} > 0$ such that $\beta_{\min} < \beta_D \lesssim h^{-1}$ and $\tau \approx h^{-1}$, then*

$$\begin{aligned} \|(u - u_h, \lambda - \lambda_h)\|_{\mathbb{V}} & \leq d_C((u, \lambda), (u_h, \lambda_h)) \\ & \lesssim h^{\zeta-1/2}|u|_{H^\zeta(\Gamma)} + h^{\xi+1/2}|\lambda|_{H^\xi(\bar{\Gamma})}, \end{aligned}$$

where $\zeta = \min(k+1, s)$ and $\xi = \min(l+1, r)$ for $\Lambda_h^l \in \{\text{DP}_h^l(\Gamma), \widetilde{\text{DP}}_h^l(\Gamma)\}$ and $\zeta = \min(2, s)$ and $\xi = \min(\frac{1}{2}, r)$ for $\Lambda_h^l = \text{DUAL}_h^0(\Gamma)$. Additionally,

$$\|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \lesssim h^{\zeta-1/2}|u|_{H^\zeta(\Gamma)} + h^{\xi+1/2}|\lambda|_{H^\xi(\bar{\Gamma})},$$

where \tilde{u} and \tilde{u}_h are the solutions in Ω defined by (2.6).

Proof. First, we observe that for all (v, μ) and (w, η) in \mathbb{W} ,

$$\|(v - w, \mu - \eta)\|_{\mathbb{V}} \leq d_C((v, \mu), (w, \eta)).$$

Using standard approximation results for $\Lambda_h^l \in \{\text{DP}_h^l(\Gamma), \widetilde{\text{DP}}_h^l(\Gamma)\}$ (see, e.g., [19, Chapter 10]) and Lemma 3.1 for $\Lambda_h^l = \text{DUAL}_h^0(\Gamma)$, we see that

$$\begin{aligned} \inf_{(v_h, \mu_h) \in \mathbb{V}_h} \|(u - v_h, \lambda - \mu_h)\|_{\mathbb{V}} & = \inf_{v_h \in P_h^k(\Gamma)} \|u - v_h\|_{H^{1/2}(\Gamma)} + \inf_{\mu_h \in \Lambda_h^l(\Gamma)} \|\lambda - \mu_h\|_{H^{-1/2}(\Gamma)} \\ & \lesssim h^{\zeta-1/2}|u|_{H^\zeta(\Gamma)} + h^{\xi+1/2}|\lambda|_{H^\xi(\bar{\Gamma})}, \\ \inf_{v_h \in P_h^k(\Gamma)} \|u - v_h\|_{\Gamma} & \lesssim h^\zeta|u|_{H^\zeta(\Gamma)}, \quad \inf_{\mu_h \in \Lambda_h^l(\Gamma)} \|\lambda - \mu_h\|_{\Gamma} \lesssim h^\xi|\lambda|_{H^\xi(\bar{\Gamma})}. \end{aligned}$$

Applying these to the definition of $\|\cdot\|_*$ gives

$$\begin{aligned} \inf_{(v_h, \mu_h) \in \mathbb{V}_h} \|(u - v_h, \lambda - \mu_h)\|_* & \lesssim h^{\zeta-1/2}|u|_{H^\zeta(\Gamma)} + h^{\xi+1/2}|\lambda|_{H^\xi(\bar{\Gamma})} \\ & \quad + \beta_D^{1/2}h^\zeta|u|_{H^\zeta(\Gamma)} + \tau^{1/2}h^\zeta|u|_{H^\zeta(\Gamma)} + \tau^{-1/2}h^\xi|\lambda|_{H^\xi(\bar{\Gamma})}. \end{aligned}$$

By means of Lemma 5.5 and the given choice of the parameters τ and β_D , we prove the first assertion. The estimate in the domain Ω follows by using the relations (2.8) and (2.10). \square

If λ is smooth enough and $k = l$, the bounds on τ can be replaced with $h \lesssim \tau \lesssim h^{-1}$ without reducing the order of convergence.

6. Numerical results. We now demonstrate the theory with a series of numerical examples. In this section, we consider the following test problem. Let $\Omega = [0, 1] \times [0, 1] \times [0, 1]$ be the unit cube, $\Gamma_C := \{(x, y, z) \in \Gamma : z = 1\}$, and $\Gamma_D := \Gamma \setminus \Gamma_C$. Let

$$(6.1a) \quad g_D = 0,$$

$$(6.1b) \quad g_C = \begin{cases} \sin(\pi x) \sin(\pi y) \sinh(\sqrt{2}\pi), & x \leq \frac{1}{2}, \\ \sin(\pi y) \sinh(\sqrt{2}\pi), & x > \frac{1}{2}, \end{cases}$$

$$(6.1c) \quad \psi_C = \begin{cases} \sqrt{2}\pi \sin(\pi x) \sin(\pi y) \cosh(\sqrt{2}\pi), & x \geq \frac{1}{2}, \\ \sqrt{2}\pi \sin(\pi y) \cosh(\sqrt{2}\pi), & x < \frac{1}{2}. \end{cases}$$

It can be shown that

$$u(x, y, z) = \sin(\pi x) \sin(\pi y) \sinh(\sqrt{2}\pi z)$$

is the solution to (1.1) with these boundary conditions.

To solve the nonlinear system (4.10), we will treat the nonlinear term explicitly. Therefore, we define

$$(6.2) \quad \mathcal{B}'_C[(u, \lambda), (v, \mu)] := \frac{1}{2} \langle \lambda, v \rangle_{\Gamma_C} + \langle \tau^{-1} \lambda - \frac{1}{2} u, \mu \rangle_{\Gamma_C}.$$

Note that \mathcal{B}'_C differs from \mathcal{B}_C only by the missing nonlinear term.

We pick initial guesses $(u_0, \lambda_0) \in \mathbb{V}_h$ and define $(u_{n+1}, \lambda_{n+1}) \in \mathbb{V}_h$, for $n \in \mathbb{N}$, to be the solution of

$$(6.3) \quad (\mathcal{A} + \mathcal{B}_D + \mathcal{B}'_C)[(u_{n+1}, \lambda_{n+1}), (v_h, \mu_h)] \\ = \mathcal{L}_C(v_h, \mu_h) - \langle [P^\tau(u_n, \lambda_n)]_+, v_h + \tau^{-1} \mu_h \rangle_{\Gamma_C} \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

This leads us to Algorithm 6.1, an iterative method for solving the contact problem.

In all of the computations in this section, we preconditioned the GMRES solver using a mass matrix preconditioner applied blockwise from the left, as described in [3].

Algorithm 6.1. Iterative algorithm for solving the contact problem.

```

Input  $(u_0, \lambda_0)$ , TOL, MAXITER
for  $n \leftarrow 0$  to MAXITER do
   $(u_{n+1}, \lambda_{n+1}) \leftarrow$  solution of (6.3), calculated using GMRES
  if  $\|(u_{n+1}, \lambda_{n+1}) - (u_n, \lambda_n)\|_{\mathbb{V}} < \text{TOL}$  then
    return  $(u_{n+1}, \lambda_{n+1})$ 
  end if
end for

```

Inspired by the parameter choices in [2], we fix $\beta_D = 0.01$ and look for suitable values of the parameter τ . Figure 2 shows how the error, number of outer iterations, and the average number of GMRES iterations inside each outer iteration change as the parameter τ is varied for both $\mathbb{V}_h = P_h^1(\Gamma) \times \text{DUAL}_h^0(\Gamma)$ (left, blue) and $\mathbb{V}_h = P_h^1(\Gamma) \times \text{DP}_h^0(\Gamma)$ (right, orange). Here, we see that the error and number of outer iterations are lowest when τ is between around 1 and 10.




FIG. 2. The dependence of the error, number of outer iterations, and the average number of GMRES iterations on τ for the problem (1.1) with boundary conditions (6.1) on the unit cube with $h = 2^{-2}$ (triangles), $h = 2^{-3.5}$ (diamonds), and $h = 2^{-5}$ (pentagons). Here we take $u_0 = \lambda_0 = 0$, $\beta_D = 0.01$, $TOL = 0.05$, and $MAXITER = 50$. On the left (blue), we take $(u_n, \lambda_n), (v_h, \mu_h) \in P_h^1(\Gamma) \times DUAL_h^0(\Gamma)$; on the right (orange), we take $(u_n, \lambda_n), (v_h, \mu_h) \in P_h^1(\Gamma) \times DP_h^0(\Gamma)$. (Figure in color online.)

Motivated by Figure 2 and the bounds in Theorem 5.6, we take $\tau = 0.5/h$, and look at the convergence as h is decreased. Figure 3 shows how the error and iteration counts vary as h is decreased when $\mathbb{V}_h = P_h^1(\Gamma) \times DUAL_h^0(\Gamma)$ (left, blue circles) and $\mathbb{V}_h = P_h^1(\Gamma) \times DP_h^0(\Gamma)$ (right, orange squares).

For $\mathbb{V}_h = P_h^1(\Gamma) \times DUAL_h^0(\Gamma)$, we observe slightly higher than the order 1 con-




FIG. 3. The error, number of outer iterations and average number of inner GMRES iteration for the problem (1.1) with boundary conditions (6.1) on the unit cube as h is reduced. Here we take $u_0 = \lambda_0 = 0$, $\beta_D = 0.01$, $\text{TOL} = 0.05$, $\text{MAXITER} = 200$, and $\tau = 0.5/h$. On the left (blue circles), we take $(u_n, \lambda_n), (v_h, \mu_h) \in P_h^1(\Gamma) \times \text{DUAL}_h^0(\Gamma)$; on the right (orange squares), we take $(u_n, \lambda_n), (v_h, \mu_h) \in P_h^1(\Gamma) \times \text{DP}_h^0(\Gamma)$. The dashed lines show order 1 convergence (left) and order 1.5 convergence (right). (Figure in color online.)

vergence predicted by Theorem 5.6. In this case, the mass matrix preconditioner is effective, as the number of GMRES iterations required inside each outer iteration is reasonably low, and only grows slowly as h is decreased. We believe that the effectiveness of the preconditioner for this choice of spaces is due to the spaces $P_h^1(\Gamma)$ and $\text{DUAL}_h^0(\Gamma)$ forming an inf-sup stable pair [18, Lemma 3.1].

When $\mathbb{V}_h = P_h^1(\Gamma) \times DP_h^0(\Gamma)$, Theorem 5.6 tells us to expect order 1.5 convergence. However, we observe a slightly lower order. This appears to be due to the ill-conditioning of this system, and the mass matrix preconditioner being ineffective, leading to an inaccurate solution when using GMRES. In this case, the spaces $P_h^1(\Gamma)$ and $DP_h^0(\Gamma)$ do not form an inf-sup stable pair, and so the mass-matrix between them is not guaranteed to be invertible leading to a less effective preconditioner.

In order to obtain order 1.5 convergence with a well-conditioned system, we could look for $(u_h, \lambda_h) \in P_h^1(\Gamma) \times DP_h^0(\Gamma)$ and test with $(v_h, \mu_h) \in DUAL_h^1(\Gamma) \times DUAL_h^0(\Gamma)$, where $DUAL_h^1(\Gamma)$ is the space of piecewise linear functions on the dual grid that forms an inf-sup stable pair with the space $DP_h^0(\Gamma)$, as defined in [4]. With this choice of spaces, we obtain the higher order convergence as in Theorem 5.6, while having stable dual pairings and hence more effective mass matrix preconditioning.

For the problems discussed in [2], we have run numerical experiments using this space pairing and observe the full order $\frac{3}{2}$ convergence in a low number of iterations. A deeper investigation of this method using these dual spaces, and the adaption of the theory to this case, warrants future work.

7. Conclusions. Based on our work in [2], we have analyzed and demonstrated the effectiveness of Nitsche type coupling methods for boundary element formulations of contact problems.

An open problem is preconditioning. While the iteration counts in the presented examples were already practically useful, for large and complex structures preconditioning is still essential. The hope is to use the properties of the Calderón projector to build effective operator preconditioning techniques for the presented Nitsche type frameworks.

Avenues of future research include looking at how this approach could be applied to problems in linear elasticity, and an extension of this method to problems involving friction.

REFERENCES

- [1] P. ALART AND A. CURNIER, *A mixed formulation for frictional contact problems prone to Newton like solution methods*, Comput. Methods Appl. Mech. Engrg., 92 (1991), pp. 353–375.
- [2] T. BETCKE, E. BURMAN, AND M. W. SCROGGS, *Boundary element methods with weakly imposed boundary conditions*, SIAM J. Sci. Comput., 41 (2019), pp. A1357–A1384, <https://doi.org/10.1137/18M119625X>.
- [3] T. BETCKE, M. W. SCROGGS, AND W. ŚMIGAJ, *Product algebras for Galerkin discretizations of boundary integral operators and their applications*, ACM Trans. Math. Software, 46 (2020), 4.
- [4] A. BUFFA AND S. H. CHRISTIANSEN, *A dual finite element complex on the barycentric refinement*, Math. Comp., 76 (2007), pp. 1743–1769.
- [5] E. BURMAN, P. HANSBO, AND M. G. LARSON, *The penalty-free Nitsche method and nonconforming finite elements for the Signorini problem*, SIAM J. Numer. Anal., 55 (2017), pp. 2523–2539, <https://doi.org/10.1137/16M107846X>.
- [6] A. CHERNOV, M. MAISCHAK, AND E. STEPHAN, *A priori error estimates for hp penalty BEM for contact problems in elasticity*, Comput. Methods Appl. Mech. Engrg., 196 (2007), pp. 3871–3880.
- [7] F. CHOULY AND P. HILD, *A Nitsche-based method for unilateral contact problems: Numerical analysis*, SIAM J. Numer. Anal., 51 (2013), pp. 1295–1307, <https://doi.org/10.1137/12088344X>.
- [8] F. CHOULY, P. HILD, AND Y. RENARD, *Symmetric and non-symmetric variants of Nitsche’s method for contact problems in elasticity: theory and numerical experiments*, Math. Comp., 84 (2015), pp. 1089–1112.
- [9] H. GIMPERLEIN, F. MEYER, C. ÖZDEMİR, AND E. P. STEPHAN, *Time domain boundary ele-*

- ments for dynamic contact problems, *Comput. Methods Appl. Mech. Engrg.*, 333 (2018), pp. 147–175.
- [10] H. HAN, *A direct boundary element method for Signorini problems*, *Math. Comp.*, 55 (1990), pp. 115–128.
- [11] H.-D. HAN, *The boundary finite element methods for Signorini problems*, in *Numerical Methods for Partial Differential Equations*, Y.-I. Zhu and B.-Y. Guo, eds., *Lecture Notes in Math.* 1297, Springer, Berlin, Heidelberg, 1987, pp. 38–49.
- [12] J. L. LIONS AND G. STAMPACCHIA, *Variational inequalities*, *Comm. Pure Appl. Math.*, 20 (1967), pp. 493–519.
- [13] M. MAISCHAK AND E. P. STEPHAN, *Adaptive hp-versions of BEM for Signorini problems*, *Appl. Numer. Math.*, 54 (2005), pp. 425–449.
- [14] J. NITSCHKE, *Über ein Variationsprinzip zur Lösung von Dirichlet-Problemen bei Verwendung von Teilräumen, die keinen Randbedingungen unterworfen sind*, *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, 36 (1971), pp. 9–15.
- [15] H. SCHMIT AND G. SCHNEIDER, *Boundary element solution of the Dirichlet-Signorini problem by a penalty method*, *Appl. Anal.*, 51 (1993), pp. 175–186.
- [16] M. W. SCROGGS, *Efficient Computation and Applications of the Calderón Projector*, Ph.D. thesis, University College London, London, UK, 2019.
- [17] W. SPANN, *On the boundary element method for the Signorini problem of the Laplacian*, *Numer. Math.*, 65 (1993), pp. 337–356.
- [18] O. STEINBACH, *On a generalized L_2 projection and some related stability estimates in Sobolev spaces*, *Numer. Math.*, 90 (2002), pp. 775–786.
- [19] O. STEINBACH, *Numerical Approximation Methods for Elliptic Boundary Value Problems*, Springer-Verlag, New York, 2008.
- [20] O. STEINBACH, *Boundary element methods for variational inequalities*, *Numer. Math.*, 126 (2014), pp. 173–197.
- [21] R. TEMAM, *Navier-Stokes Equations: Theory and Numerical Analysis*, *Studies in Mathematics and its Applications*, Vol. 2, North-Holland, Amsterdam, New York, Oxford, 1977.
- [22] S. ZHANG AND X. LI, *An augmented Lagrangian method for the Signorini boundary value problem with BEM*, *Bound. Value Probl.*, 2016 (2016), 62.
- [23] S. ZHANG AND J. ZHU, *A projection iterative algorithm boundary element method for the Signorini problem*, *Eng. Anal. Bound. Elem.*, 37 (2013), pp. 176–181.